Mon. Not. R. Astron. Soc. **000**, 1–?? (2004) Printed 2 February 2008 (MN LaTeX style file v2.2)

arXiv:astro-ph/0407023v1 1 Jul 2004

Revisiting the Magnification of Type Ia Supernovae with SDSS

Brice Ménard and Neal Dalal*
Institute for Advanced Study, Einstein Drive, Princeton NJ 08540 USA

2 February 2008

ABSTRACT
We cross-correlate the sample of Type Ia supernovae from Riess et al. (2004) with the SDSS DR2 photometric galaxy catalogue. In contrast to recent work, we find no detectable correlation between supernova magnitude and galaxy overdensity on scales ranging between $1 - 10'$. Our results are in accord with theoretical expectations for gravitational lensing of supernovae by large-scale structure. Future supernova surveys like SNAP will be capable of detecting unambiguously the predicted lensing signal.

Key words:

1 INTRODUCTION

Weak gravitational lensing by large-scale structure can both distort and magnify the images of background sources. The former effect, cosmic shear, has been detected (see Réfrégier 2003 for a review) and holds great promise for the study of matter fluctuations on large scales. The latter effect, cosmic magnification, has yet to be detected convincingly but has been theoretically investigated in detail (Jain et al. 2003; Ménard et al. 2003; Takada & Hamana 2003). In particular, a number of authors have studied magnification of Type Ia supernova, which have been employed as standard candles to measure the expansion history of the universe (e.g. Tonry et al. 2003; Knop et al. 2003). The RMS magnification of supernovae is expected to be at the $\sim 1\%$ level on arcminute scales, with the largest magnifications occurring near the centers of massive halos.

On scales larger than arcminutes, only modest shear and magnification at the sub-percent level are expected. The theoretically expected magnitude of shear fluctuations has been confirmed observationally in cosmic shear surveys. Accordingly, it would be extremely surprising if observations contradicted the expected level of cosmic magnification, since statistical shear and magnification are related in a simple way. In principle, standard candles like Type Ia supernovae should offer a means of measuring the magnification caused by large scale structure, for example by cross-correlating supernova brightness with foreground galaxy overdensity. In practice, all but the very highest redshift supernovae (which suffer the most lensing) are not expected to provide useful measurements of cosmic magnification, since even Type Ia SNe are not perfect standard candles. The observed dispersion in supernova magnitudes is roughly $\delta m \sim 20\%$, whereas fluctuations of only $\lesssim 1\%$ are expected from lensing (Ménard et al. 2003; Takada & Hamana 2003). Even if magnification and galaxy overdensity were perfectly correlated, we would expect to see only a weak correlation between supernova brightness and galaxy overdensity, $r = \sqrt{\langle\delta\mu^2\rangle/\langle\delta m^2\rangle} \lesssim 0.05$. Therefore a very large sample of SNIa would be needed to observe such an effect.

Recently, Williams & Song (2004) have reported a highly significant ($> 99\%$ confidence) detection of correlation between the magnitude of 55 supernovae and foreground galaxy overdensity on $10'$ scales, corresponding to magnifications at the $\sim 10\%$ level. Wang (2004) has claimed further evidence for gravitational magnification of the Riess et al. supernovae. If confirmed, these results would confound much of the conventional wisdom on magnification effects and would call into question previous cosmic shear measurements which detected much lower shear variance on these scales. Additionally, if such a correlation exists, it is possible to correct for it and therefore significantly reduce the dispersion in supernova luminosities: the magnitude variance can be reduced by a factor $1-r^2$ (Dalal et al. 2003). This would be especially important for modern supernova surveys, which can calibrate supernova magnitudes nearly to 0.1 mag.

In this paper, we revisit this issue by cross-correlating the most recent sample of cosmological Type Ia SNe compiled by Riess et al. (2004), with low redshift galaxies in the Sloan Digital Sky Survey (SDSS) photometric catalogue. The plan of the paper is as follows. In section 2, we briefly review theoretical predictions for cosmic magnification on small scales. Next, we describe the supernova sample and galaxy catalogue employed in our analysis. Results are presented in §4, and our conclusions are discussed in §5.

---

* Hubble Fellow

2 COSMIC MAGNIFICATION

The large scale distribution of matter in the Universe can give rise to measurable gravitational lensing effects (shear and magnification) on background objects. The magnification factor μ depends on whether matter along the line of sight is preferentially over- or underdense compared to the mean.

The local properties of the gravitational lens mapping are characterised by the convergence κ, which is proportional to the surface mass density projected along the line-of-sight, and the shear γ, which is related to the gravitational tidal field of the lensing mass distribution. The SDSS population of galaxies with $m_r < 21$ has a median redshift near 0.3 (e.g. Dodelson et al. 2002), so that angular scales of arcminutes correspond to length scales of Mpc. On these scales, the convergence and shear fluctuations are small, $|\kappa|, |\gamma| \ll 1$. To leading order the magnification is approximately

$$\mu = 1 + 2\kappa + \mathcal{O}(\kappa^2, |\gamma|^2)\ , \tag{1}$$

see Ménard et al. (2003) for higher order corrections. Observable effects are due to departures from the mean value of the magnification. It is therefore convenient to use the overmagnification $\delta\mu = \mu - 1$. Then, the cross-correlation between $\delta\mu$ and the foreground matter overdensity traced by galaxies at an angular separation θ reads:

$$\begin{aligned}\langle \delta\mu\, \delta_{\rm gal}\rangle(\theta) &\approx 2\, \langle \kappa \times \delta_{\rm DM}\rangle(\theta) \\ &\approx \frac{3}{2}\, \Omega_0 \left(\frac{H_0}{c}\right)^2 \int \frac{{\rm d}w}{a(w)}\, \frac{D_d\, D_{ds}}{D_s}\, W_{\rm gal}(w) \\ &\quad \times b_{\rm gal} \int \frac{s{\rm d}s}{2\pi} P_\delta\left(\frac{s}{f_k(w)}, w\right) {\rm J}_0(s\,\theta)\ . \end{aligned} \tag{2}$$

In this expression, w is the comoving distance along the line-of-sight, $D_{d,s,ds}$ are angular diameter distances to lens, source, and from lens to source respectively, $W_{\rm gal}$ is the normalised distribution of galaxies along the line-of-sight, a is the scale factor and P_δ is the matter power spectrum, calculated using $\Gamma = 0.19$ and the fitting function of Peacock & Dodds (1996), and we assume that the linear bias $b_{\rm gal} = 1$ for the galaxies and scales of interest (Verde et al. 2002). In principle, the bias of the detected galaxies should vary with redshift, but because the galaxy distribution is dominated by a narrow redshift interval, the use of a constant bias seems reasonable. If instead of the two-point correlation, we wanted the cross-correlation with galaxy overdensity smoothed on scale θ we would replace ${\rm J}_0(s\theta)$ by $2{\rm J}_1(s\theta)/s\theta$.

Figure 1 shows the amplitude of this cross-correlation between $\delta\mu$ and $\delta_{\rm gal}$ (Eq. 2). The solid line shows $\langle\delta\mu\, \delta_{\rm gal}\rangle$ computed for different angular separations and the dashed line shows the integrated correlation, i.e. the correlation between supernova magnitude and the number of galaxies within radius θ. For estimating these quantities we have used a galaxy redshift distribution peaking at $z = 0.3$ and a WMAP cosmology with $\Omega_m = 0.27$, $\Omega_\Lambda = 0.73$, $n = 1$, and $\sigma_8 = 0.9$ (Spergel et al. 2003). Black lines have been computed for supernova redshifts of $z_s = 0.6$ (the median redshift of our sample) and gray lines for $z_s = 2$. As can be seen in this figure, magnifications of $\lesssim 1\%$ can be generated by structures on scales of a few arcminutes or larger.



Figure 1. Cross-correlation between gravitational magnification and galaxy overdensity as a function of angular scale. A linear bias of unity has been used in this calculation.

As an aside, it is perhaps worth noting that high redshift sources are expected to be more strongly perturbed by lensing fluctuations, both because the lensing efficiency increases with redshift, and because more distant sources are subject to lensing from structures along a greater path length. For example, sources at $z = 2$ are typically magnified by $\delta\mu \sim 2\%$ by structures on scales of $1'$. This has ramifications for pencil-beam surveys attempting to constrain dark energy with supernovae at high redshift, $z \sim 1.5$. For example, Riess et al. (2004) report the discovery of 16 SNIa in the GOODS field, over an area of 0.1 sq. degrees. Lensing fluctuations are significantly correlated across this narrow field. If these high redshift supernovae played an important role in the constraints derived by Riess et al., then their derived dark energy parameters could be significantly biased, for example by 5% in w. Fortunately, the 16 high redshift GOODS supernovae were negligible within the larger sample of 177 SNe reported by Riess et al., implying that their dark energy constraints were not appreciably biased. Future wide area surveys, like the SNAP survey which expects to cover 20 sq. degrees, should also be undisturbed by correlated structure across their fields.

3 THE DATA

We attempt to detect a correlation between supernova magnitude and galaxy overdensity using following data sets:

- For supernovae, we use the "gold" sample of 156 SNIa compiled by Riess et al. (2004). For each supernova, the redshift, (dust-corrected) distance modulus, and extinction correction have been provided by these authors. The SN redshifts range from $z = 0.014$ to 1.75 with an average value of ≈ 0.6.
- For foreground galaxies, we use the second data release (DR2) of the Sloan Digital Sky Survey (SDSS). As shown by Scranton et al. (2002) this dataset provides an efficient star/galaxy separation for objects with $r < 21$ and a homogeneous density by requiring a seeing value greater than $1.5''$.

Of the 156 SNe listed by Riess et al., 55 fall within the SDSS DR2 footprint. Given the magnitude cut mentioned above, the average galaxy density is roughly 1 arcmin^{-2}. The list of SNIa that we use is given in table 1.

Williams & Song used magnitude cuts in each field in order to restrict their galaxy counts to low redshifts objects and maximize the signal. In our case, the bulk of SDSS photometric galaxies are at redshifts $z \lesssim 0.3$, while the median SN redshift in our sample is $z_{\rm med} = 0.6$. Accordingly, the number of galaxies that will overlap the supernova redshift distribution of our sample will be largely negligible.

4 ANALYSIS

For each of the 55 SNIa we first count the number of neighbours within an angular radius θ. The correlation between this number of neighbours and the absolute SNIa magnitude is shown in Fig. 2. The left panel shows, as a function of supernova magnitude, the distribution of galaxies located within $10'$ from the SNIa position, i.e. the angular scale where Williams & Song (2004) found the strongest correlation. The horizontal error bars correspond to the measurement error in the SNIa magnitudes quoted by Riess et al. (2004), and the vertical errors correspond to the Poisson noise of the galaxy counts. There is no apparent correlation in this scatter plot; this will be quantified below. The right panel shows the same correlation but using stars instead of galaxies and allows us to check for possible systematics.

For each angular bin, we have computed the cross-correlation coefficient

$$r = \frac{\left\langle (\Delta N - \bar{N})\,(\Delta M - \bar{M}) \right\rangle}{\sqrt{\langle (\Delta N - \bar{N})^2 \rangle \langle (\Delta M - \bar{M})^2 \rangle}}\,, \qquad (3)$$

where N is the number of neighbours (galaxies or stars) within a given angular separation from the supernova and M is the SN absolute magnitude. Neglecting the measurement uncertainties in each object, and accounting only for the scatter in the population, the error in the cross-correlation coefficient reads

$$\sigma_r = \frac{1 - r^2}{\sqrt{N_{\rm SN}}}\,, \qquad (4)$$

where $N_{\rm SN}$ is the number of SNIa. For cells of 10 $'$radius, we find $r = 0.03 \pm 0.13$ and $r = -0.04 \pm 0.26$ for galaxies and stars respectively. The result obtained by Williams & Song (2004) is therefore not reproduced in our analysis: no significant correlation is detected.

In Fig. 3 we plot the cross-correlation coefficient r as a function of θ. In this figure, we have counted galaxies within an annulus centred on each θ, so that the plotted errors are uncorrelated. For both galaxies and stars, no significant correlation has been detected on any scale.



Figure 3. Cross-correlation coefficient r between SN magnitude and galaxy counts (solid) or star counts (dashed), as a function of radius. The overplotted curves show the theoretical prediction for source redshifts of 0.6 and 2.

5 DISCUSSION

Using the "gold" sample of type Ia supernovae given by Riess et al. (2004) and the galaxy distribution measured by SDSS DR2, we have investigated correlations between the brightness of the supernovae and the density of foreground galaxies, on scales of 1-10$'$. Our analysis does not reproduce the recent results of Williams & Song (2004), who detected a cross-correlation at high significance (> 99% confidence) on $10'$ scales. Our result is in line with theoretical predictions: the $\lesssim 1\%$ rms magnification caused by large-scale structure on arcminute scales is negligible compared to the observed $\sim 20\%$ dispersion in supernova magnitudes.

The number of SNe used by Williams & Song closely matches the number used in our analysis. The main difference between the two analyses is that Williams & Song derive galaxy counts from the APM catalogue, whereas we have used counts from the SDSS photometric catalogue. While the APM catalogue is based on scans of photographic plates taken with the UK Schmidt telescope, the SDSS uses modern CCDs with high-quality imaging reaching down to much deeper magnitudes, providing a higher galaxy density. Therefore we expect the SDSS to be more suitable for investigating weak correlations.

Thus there is no evidence for a significant correlation between supernova brightness and foreground galaxy overdensity. Such a correlation would be expected for a sample of weakly lensed supernovae. In this light, the recently claimed detection of significant weak lensing in the Riess et al. (2004) sample by Wang (2004) is somewhat surprising. Further examination of the data presented by Wang indicates that the effect appears to be dominated by a few outliers in the magnitude distribution, which Wang ascribes to magnification factors ~ 2, i.e. strong lensing. This would imply a surprisingly high strong lensing rate ($\sim 5\%$) among high z supernovae, inconsistent with the strong lensing rate ($\sim 10^{-3}$) of distant quasars (Browne et al. 2003).

Lastly, we note that the theoretically expected SN-galaxy correlation (solid curves in figure 3) would be detectable with larger samples of supernovae with smaller flux errors. Although the signal is not currently detectable using the Riess et al. (2004) supernovae and SDSS galaxies,



Figure 2. Scatter plot of residual magnitude vs. number of galaxies (left) or stars (right) within 10′.

future datasets (e.g. SNAP) should unambiguously exhibit the lensing signal on small scales.

ACKNOWLEDGMENTS

BM is supported by a fellowship of the F. Gould foundation. ND is supported by NASA through Hubble Fellowship grant #HST-HF-01148.01-A, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555.

REFERENCES

Blandford, R. D., Saust, A. B., Brainerd, T. G. and Villumsen, J. V., 1991, MNRAS, 251, 600

Browne, I. W. A., et al. 2003, MNRAS, 341, 13

Dalal N., Holz D. E., Chen X., and Frieman J. A. 2003, ApJ, 585, L11

Dodelson, S., et al. 2002, ApJ, 572, 140

Jain, B. and Scranton, R. and Sheth, R. K. 2003, MNRAS, 345, 62

Knop, R. A., et al. 2003, ApJ, 598, 102

Ménard, B., Hamana, T., Bartelmann, M., & Yoshida, N. 2003, A&A, 403, 817

Peacock, J. A. & Dodds, S. J. 1996, MNRAS, 280, L19

Refregier, A. 2003, ARA&A, 41, 645

Riess, A. G., Strolger, L., Tonry, J., Casertano, S., Ferguson, H. C., Mobasher, B., Challis, P., Filippenko, A. V., Jha, S., Li, W., Chornock, R., Kirshner, R. P., Leibundgut, B., Dickinson, M., Livio, M., Giavalisco, M., Steidel, C. C., Benitez, N., & Tsvetanov, Z. 2004, accepted to ApJ, astro-ph/0402512

Scranton, R., et al. 2002, ApJ, 579, 48

Spergel, D. N., Verde, L., Peiris, H. V., Komatsu, E., Nolta, M. R., Bennett, C. L., Halpern, M., Hinshaw, G., Jarosik, N., Kogut, A., Limon, M., Meyer, S. S., Page, L., Tucker, G. S., Weiland, J. L., Wollack, E., & Wright, E. L. 2003, ApJS, 148, 175

Takada, M. and Hamana, T. 2003, MNRAS, 346, 949

Tonry, J. L., et al. 2003, ApJ, 594, 1

Verde, L., et al. 2002, MNRAS, 335, 432

Wang, Y. 2004, astro-ph/0406635

Williams, L. L. R. and Song, J. 2004, accepted to MNRAS, astro-ph/0403680

Table 1. Supernovae used in our analysis. Here, μ_0 and σ are the distance modulus and its error. All quantities are taken from the larger "gold" sample of Riess et al. (2004).

SN	RA	DEC	redshift	μ_0	σ	host A_v
1999dk	01 31.5	+14 17	0.0141	34.43	0.26	0.20
1993ae	01 29.8	-01 59	0.0180	34.29	0.23	0.00
1994M	12 31.2	+00 36	0.0244	35.09	0.20	0.23
1994Q	16 49.9	+40 26	0.0290	35.70	0.19	0.33
1998cs	16 30.7	+41 13	0.0327	36.08	0.19	-0.03
1994T	13 19.0	-02 09	0.0360	36.01	0.20	0.09
1997I	04 59.6	-03 09	0.172	39.79	0.18	—
1999fw	23 31.9	+00 10	0.278	41.00	0.41	0.26
1997bj	10 42.4	+00 02	0.334	40.92	0.30	0.34
1996K	08 24.7	-00 21	0.380	42.02	0.22	0.02
1995ba	08 19.1	+07 43	0.3880	42.07	0.19	—
1997am	10 57.5	-03 14	0.416	42.10	0.19	0.00
1997bh	13 44.6	-00 20	0.420	41.76	0.23	0.60
1997Q	10 56.9	-03 59	0.430	41.99	0.18	—
1998ba	13 43.6	+02 20	0.430	42.36	0.25	—
1997aw	10 23.5	+04 07	0.440	42.57	0.40	0.80
1997ai	10 49.0	+00 32	0.450	42.10	0.23	—
1999ff	02 33.9	+00 33	0.455	42.29	0.28	0.19
1997P	10 55.9	-03 57	0.472	42.46	0.19	—
2002dc	12 36.8	+62 13	0.475	42.14	0.19	0.23
1995ay	03 01.1	+00 21	0.480	42.37	0.20	—
1996ci	13 45.9	+02 27	0.495	42.25	0.19	—
1997cj	12 37.1	+62 26	0.500	42.74	0.20	0.15
1999U	09 26.7	-05 38	0.500	42.75	0.19	0.04
1997as	08 24.2	-00 48	0.508	41.64	0.35	0.85
1997bb	12 29.0	+00 09	0.518	42.83	0.30	0.11
2001iy	10 52.4	+57 17	0.570	42.88	0.31	-0.04
1996I	12 00.7	-00 16	0.570	42.81	0.25	0.14
1997af	08 23.9	+04 09	0.579	42.86	0.19	—
1995ax	02 26.4	+00 49	0.615	42.85	0.23	—
1996H	12 28.9	-00 05	0.620	43.11	0.30	0.09
1998M	11 33.7	+04 05	0.630	42.62	0.24	0.75
2003be	12 36.4	+62 07	0.64	43.07	0.21	0.23
1997R	10 57.3	-03 55	0.657	43.27	0.20	—
2003bd	12 37.4	+62 13	0.67	43.19	0.28	0.27
2001ix	10 52.3	+57 07	0.710	43.05	0.32	0.53
1998bi	13 47.7	+02 21	0.740	43.35	0.30	—
1997ez	08 21.6	+03 25	0.778	43.81	0.35	—
2001hx	08 49.4	+44 02	0.798	43.88	0.31	0.31
2001hy	08 49.8	+44 15	0.811	43.97	0.35	0.03
1999fj	02 28.4	+00 39	0.815	43.76	0.33	0.23
2001jf	02 28.1	+00 27	0.815	44.09	0.28	0.23
1996cl	10 57.0	-03 38	0.828	43.96	0.46	—
1997ap	13 47.2	+02 24	0.830	43.85	0.19	—
2003eq	12 37.8	+62 14	0.839	43.86	0.22	0.22
2001jh	02 29.0	+00 21	0.884	44.23	0.19	-0.01
2003eb	12 37.3	+62 14	0.899	43.64	0.25	0.26
2003lv	12 37.5	+62 11	0.94	43.87	0.20	0.15
2002dd	12 36.9	+62 13	0.95	44.06	0.26	0.24
2003es	12 36.9	+62 13	0.954	44.28	0.31	0.07
2002ki	12 37.5	+62 21	1.140	44.84	0.30	0.09
1999fv	23 30.6	+00 17	1.19	44.19	0.34	0.24
2003az	12 37.3	+62 19	1.265	45.20	0.20	0.25
2003dy	12 37.2	+62 11	1.340	45.05	0.25	0.54
1997ff	12 36.7	+62 13	1.755	45.53	0.35	0.00